FRONT 55mm

BACK/PEAL 55mm

BACK 55mm



MITES I WHITEFLY I CATEPILLARS I APHIOS IBLACK BEETLE I MEALYBIOS SOFT MOULD I CITRIS LEAF MINER! BRONZE BEETLE I CRASS GRIB BEETLE THERS I SCALE I LEAF ROLLER! PSYLLID! POTATO BETLE HOW TO PRAPIRE: experience of the property of the property of the property of the property of the leaf of covering and suding include it grade. Consistent applications are necessary during gest activity. Mix will be before spropsy, hold: There could be some species with these pest groups that may not be controlled.

COMPATIBLITY: cen-neem? Organic bottomal blescaleds is compatible with experient and preparing product for use, were number glows. Wish hands after use. After each day's use week glows. Down of allow themic continenes or garpy to get into drains, sevens, stream or ponds. Do Not! feet treated grass clippage to levelstock.

PLAIT SAFETY: Trais on resemble plants such as ferns, potentiestic and Aftern widels have recutted in leaf durings. The property of t

rubbish.

FIRST AID: If poisoning occurs, contact a doctor or
National Poisons Centre 0800 POISON (0800 764 766).

FOR EMERGENCIES ONLY CALL: 0800 220 770
(ALL HOURS).

6 Magnification Barcode

HOW TO LISE

lant/Situation	Pest	Rate	How to Apply
Vegetables, Fruit, Ornamental plants	Caterpillars Sooty mould	Mix 10mL per 5L water	Apply when pests appear. Re-apply every 7 days while pests are present. Good coverage is important so apply spray to upper and underside of leaves to point of nu-off. Note plant safely CAUTION information. Caterpilliars: Use of an emulsified botanical spray additive. e.g. caterpilliars: Use of an emulsified botanical spray additive. e.g. eco-off is recommended. Note: There may be some species within this pest group that may not be controlled. Sooty mould: Spray insects directly above and around or within the soly mould affected areas. Meathylugs: use the higher rate when seet furnithers are high.
	Aphids, Leaf miner, Leaf Roller, Mittes, Potato Beetle, Scale, Thrips, Whitefly Bronze Beetle, Mealybugs, Psyllid	Mix 15-25mL per 5L water Mix 25mL per 5L water	
Home garden lawns, plant pots and garden beds (ornamentals only)	Grass Grub Beetle	Mix 20mL per 5L water to treat 10 square metres	Apply at the first sign of pest infestation and before pest populations build up to damaging levels. Ensure thorough coverage of treated areas. Apply using a watering can.

Plant/ Situation	Pest	Rate	How to Apply
Home garden lawns, plant pots and garden beds (ornamentals only)	Black Beetle larvae	Mix 10mL per 1L water	Apply as a soil drench using a watering can. For Black Beetle, the optimum application time is peak egg hatch which occurs between mid Spring to mid Summer. Ensure thorough coverage of treated areas.
Ornamental citrus	Citrus leaf miner	Mix 25mL per 10L water	Begin applying when new flush growth is 4 cm long. Apply every 5 to 14 days with presence of new flush growth. Ensure that new flush leaves are sprayed on top and underneath.

BACK

55mm

WITHHOLDING PERIOD (WHP): 1 DAY
IT IS AN OFFENCE TO USE THIS PRODUCT ON ANIMALS.
STORAGE & DISPOSAL: Store in the FIRST AID: If poisoning occurs, occurs of the property of the period product of the

The OCP drop device is a registered trade mark and **eco-**neem and **eco-**oil are trade marks of Organic Crop Protectants. NeemAzal - T/S is a registered trade mark of Andreas Welte